Table I.—Mean monthly data from January 1872, to December 1886.

					8	un s	pote.	•								Me	gnet	tic de	clina	tion	•			Range of air pressure (inches),								,		
Year.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	Јаве.	July,	Aug.	Sapt.	g.	Dec.
872 873 874 875 876	43 30 7	60 54 32 11 8	44 50 23 17	50 38 16 16	54 23 21 6 3	55 22 19 12 2	52 33 35 8 9	42 34 32 8 4		52 23 18 7 7	56 39 14 9 5	42 25 16 5 4	10.5 10.1 9.3 5.9 7.1	9.5 9.1 8.9 6.1 6.0	10.1 9.9 8.0 5.6 5.2	10.9 10.3 7.8 5.5 5.4	8.4 8.2 6.2 5.0	7.3 5.7 7.1	1.0 I 8.7 7.9 5.7 7.4	3.6 7.5 5.7	8.2 7 8.2 6	.9 10. 8 8. 9 7. 1 6.	2 8.4 0 6.6 4 7.0	1.05	1.08	I.07 I.12 I.04 I.00 I.12	.89 •74 •83 .69	.74 .72 .64 .87	.56 •55 •50 •53	.42 .43 .47 .45	·39 ·42 ·45 ·53 ·39	.61 .60 •45 •70 •55	.58 .77 I .77 I .75	90 .92 08 .99 16 .91 99 .77 85 .94
877 878 879 380 38 t	0 12 18	4 6 0 6 20	8 6 0 2 32	9 1 2 3 3 6	11 4 1 12 20	2 4 2 15 34	1 0 1 12 34	2 1 4 23 36	0	5	6 0 5 16 20	2 0 I II 2I	6.8 5.7 6.5 6.8 7.6	5.7 5.8 6.3 8.2	5.3 4.7 5.5 6.0 7.2	4.3 4.9 4.3 6.3 6.7	5.2 4.8 5.7 6.4 7.5	5.7 6.3 6.7 7.1 9.4	6.0 5.6 6.3 7.4 8.5	5.9 5.4 5.5 3.6	1.5 5.6 7.6 7.8	.7 6. .1 5. .8 6. .9 7. .4 7.	8 7.4 7 8.6	.97 .88 .94 1.04	.96 1.08 .88 1.18	1.00 .81 .86 .97 .81	.78 •78	t			.43 .34 .59 .51	.51 .57 .68 .62 .61	.64 .77 .82 .87 .76	73 .91 73 .91 83 .80 00 1.14 88 .82
882	56 26	32 20 35 51 14	26 15 48 42 58	57 36 61 46 32	40 7 45 73 41	29 32 24 100 28	27 49 31 62 28	16 22 32 43 14	27 27 39 36 25	27 36 28 40 7	28 33 16 27 0	12 38 25 15 [0]	7.8 7.9 8.0 8.2	8.3 8.4 8.6 6.5	7.8 8.2 8.2 6.2	7.5 7.7 8.8 5.9	9.6 7.8 8.5 7.8	7.2 9.1 9.9 9.8	6.8 9.4 7.8 9.0	6 8 5.0 8 7.0 8	3.1 9 3.5 9 3.3 8 7.3 7	.3 8. .5 8. .5 7. .9 7.	9 8.2 8 8.0 5 8.2 7.2	1.00 .99 1.03 1.19 1.02		1.01 1.04 .93 .81	.82 .95 .65 .72	.68 .70 .58 .62	.63 .50 .52 .52	.44 .55 .41 .40 .57	.44 .44 .50 .51	• 57	.83 I .60	81 1.29 03 .90 92 1.09 75 1.17 00 .99
Сог				• • • • •		•••••		·····	· ····	· ····	•••••			•••••		····· ·	•••• •	·····]					24	26	—.2 0	—.o5	+.12	+.24	+-33	+.31	20	.02 —	21 2
	Meau temperature (degrees).							Clouds (tenths).						Precipitation (inches).																				
Year.	Jan.	Feb.	Apr.	May.	June.	Aug.	Sept.	Oct.	Nov.	Jec.	Ę	Mar	Apr.	May.	June.	July.	Δng.	Sept.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	. ;	May.	Јипе.	July.	Aug.	Sept.	Oct.	Nov.	Dec.
72 73 74 75	33 30 38 29 42	38 4 38 4 39 4 33 4 42 4	2 59 6 55 7 51 4 54 3 58	68 66 70 67 68	76 77 77 74 74	79 79 78 79 80 79 79 73	8 71 8 68 9 70 8 67 68	58 56 59 55 59	42 45 47 46 44	31 5 40 5 40 5 45 5 28 5	·3 5 ·5 5 ·9 6 ·9 5	3 5 1 4 1 5 6 6 8 5	4 5. 6 5. 6 5. 9 5.	0 5. 1 5. 4 4. 7 4. 0 5.	7 5. 1 4. 4 5. 2 4.	3 4.9	4.3	4.5 5.8 4.9 4.8 4.7	3.6 3.7 3.8 4.2 3.6	5.9 5.7 5.9 6.0	6.3 5.9 5.9 5.4 4.8	13.00 29.21 20.91 26.91 39.89	18.1: 25.8: 24.0: 31.7: 24.2:	30.1 8 18.8 5 38.6 7 47.8 5 56.0	0 33. 0 27. 6 60. 3 26. 8 28.	83 41 18 54 62 11 71 22 35 40	2.09	27.25 38.23 36.51 40.18 29.96	28.84 22.69 32.06 42.38 28.62	24.1	19.0 34.1 24.9	2 22.6 9 8.2 9 14.4	18.6 2 22.6 1 31.2	8 20.59 3 17.5
	34 38	44 4	57 6 62	67 67	75 74 75 75 75 77	79 77 81 79 80 75	7 70 9 69 5 67 7 67 7 73	60 59 64 57 63	45 49 50 38	48 5 33 5 39 4 33 5 45 6	.6 5 .2 5 .8 5					1 4.4 0 3.7 3 4.2 5 4.1 1 3.5	3.6 3.7 4.4 4.5 3.6	4.9 3.6 3.6 4.8 3.2	5.3 4.2 3.9 4.8 5.6	5.5 4.0 4.7 6.2 5.2	5.4 5.8 6.3 6.2 5.2	18.35 20.34 21.58 22.07 28.63	6.9 16.4 16.7 29.0 35.3	26.5 22.8 3 12.9 38.7 3 17.1	39. 8 38. 8 26. 2 32. 8 19.	50 17 24 32 71 32 52 32 48 22	2.85 2.86 2.16 2.47	35.89 22.97	31.61 34.82 16.13	35.8	33.9 17.7. 14.6 34.7 35.8	12.0 2.25.0	9 29.4	28.16 2 30.82 9 21.02
78 79 80	38 34 48 29	44 4 44 5 37 5 44 4 36 4	58 59 55	67 67 67 71 72	75 77	80 7: 77 7 81 8	73	63	47	45 0	. 4 3	9 4	1 -				1	1 .	اء د	5.7	5.8	41.96	43.10	5 27 2	0 26	arl .6	1.07	26 24		! .			1	1 .
377 378 379 380 381 382 383 383 384 883	l í	37 5 44 4 36 4 48 5 38 4 41 4 32 4 37 4	59 57	1	75 74		73 5 69 5 68 5 74 5 69 7 71	63 64 60 63 56 61	59 50 48 48 45	37 6 42 5 37 5 39 4 32 6	.6 5 .8 5 .3 6 .4 4	3 5 7 5 0 5 6 4 7 6	2 5. 1 5. 9 6. 9 5.	0 5.3 5.0 7 4.9 3 4.9	5.0 5.1 5.4 9 4.9	5.3 4.3 2 4.6 4.5 3.4	3.5 3.7 4.3 3.9	3.8 3.5 4.0 5.1 4.7	5.6 3.6 4.2 2.8	4.6 3.8 5.0 4.9	5.3 6.9 4.7 5.0	28.59 22.55 33.15 30.83	35.6 31.7 11.9	33.4 33.4 5 14.2 8 27.5	43. 2 30. 8 27. 3 32.	05 28 12 35 60 23 72 22	.44 5.68 3.64	42.50 33.11 25.84 43.07	28.97 29.89 28.48 14.88	15.14	32.8	23.4	35.1 B 14.5 9 16.1	0 18.12 1 25.29 4 51.50 9 17.79 2 13.73

 ${\bf Table~II.--Means~of~five~consecutive~months.}$

	Sun spots.										М	agno	tic d	oclin	atio	n.				Range of air pressure (inches)																
Year.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.
1872 1873 1874 1875 1876	49 30 13	45 25 13	. 50 42 24 11 7	53 37 22 13 6	51 33 23 12 7	51 30 25 10 4	5 ² 27 24 7 5	52 27 23 7 6	52 30 22 7 6	50 29 19 6 5	50 28 14 6 7	49 30 13 7	10.0 9.7 8.6 6.2 6.3	10.1 9.8 8.5 5.9 6.1	10.3 9.6 8.4 5.9 5.7	10.4 9.3 8.0 5.8 5.7	10.7 9.2 7.8 5.7	11.1 8.9 7.7 6.0 6.3	11.1 8.5 7.8 6.3 6.4	10.8 8.4 7.6 6.3 6.8	10.6 8.3 7.5 6.4 6.6	10.3 8.2 7.2 6.7 6.5	8.4 6.9 6.7	6.5	.81	.78 .83 .75 .76	.80 .81 .82 .60	.83 .80 .79 .82 .78	.82 .80 .78 .80	.79 .76 .77 .60 .73	.78 .78 .74 .86 .74	.73 .77 .79 .81	.71 .79 .79 .81	.70 .79 .77 .77	.72 .83 .75 .76	.72 .84 .81 .74 .75
1877	4 0 5	8 3 0 6 23	9 4 1 8 25	7 4 1 9 28	6 3 1 10 31	5 2 2 14 32	4 2 2 19 30	3 21 30	3 21	3 21	3 0 5 20 21	3 0 6 17 22	6.2 5.8 6.0 6.5 7.6	5.8 5.5 5.8 6.5 7.4	5.2 5.6 6.4 7.4	7.8	5.3 5.7 6.6 7.9	5.4	5.3	5.4	5.2	5.5	5.9 5.7 6.3 7.7 8.3	5.9 6.1	.72 .09 .63 .73	.79 .73 .67 .78 .75	.79 .71 .69 .82	.78 .72 .73 .85 .67	.79 .71 .76 .79 .67	.78 .72 .81 .80 .69	.73 .73 .84 .79 .72	.72 .75 .85 .80 .74	.68 .70 .80 .79	.68 .69 .78 .85	.68 .68 .74 .85 .73	.70 .65 .75 .83 .72
1882 1883 1884 1885 1886	17 42 32	31 19 48 38 31	34 18 49 48 37	37 22 43 62 35	36 28 42 64 37	34 29 39 64 29	28 27 34 63 27	25 33 31 56 20	33 29 42	31 28 32	38 27 33	20 40 29 27	8.3 8.3	8.1	8.0	8.1 8.2 8.8 7.2	8.4	8.4	7.9 8.6 8.3 8.4	8.2	7.5	8.2 8.6 7.9 7.7	8.1	7 8	.73 .81 .75 .76	.75 .87 .74 .75 .81	.78 .81 .74 .73 .76	.81 .82 .73 .69 .76	.80 .84 .74 .70 .77	.79 .82 .76 .74 .78	.78 .79 .75 .76 .79	.75 .79 .73 .75 .83	.69 .80 .73 .71 .84	.75 .77 .75 .75 .82	.75 .77 .77 .75	.76 .77 .76 .76
		3	Iean	tenı	perat	ure	(degi	rees.))					Cı	ouds	(ten	tbe.)										F	recip	itatic	n (inc	ches.)					
Year.	Jan.	Feb.	Apr.	May.	June.	Aug.	Sept.	Oct.	Nov.	Jan	Feb.	Mar.	Apr.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.	Jan		Feb.	Har.	Apr.	. ;	ILB.	June.	July.	Aug.	Sept.	;	i det	Nov.	Dec.
1872 1873 1874 1875	54 58 55	54 57 54 60	56 50 55 59 57 53 58 59 53	57 56 57 56 56 56	58 57 57 55 55	58 5 57 5 59 5 55 5 57 5	8 57 7 56 8 58 5 55 6 56	55 57 58 57 54	54 57 57 59 54	54 58 5. 56 5. 60 5. 55 4.	0 4. 2 5. 3 5. 9 4.	8 4. 3 5. 7 4.	9 5.0	4.9 5.2 4.7 5.1	4.5 5.3 4.7 5.0 5.1	5.0 5.0 4.7 5.2	4.8 5.2 4.7 5.2 5.0	5.0 5.1 4.9 5.3 5.1	5.0 5.1 5.0 5.1 4.9	5.2	5.0	24. 26.	57 01 63	25.11 32.89 30.10	29.4 29.3	0 29. 9 30. 3 30.	87 28 61 28 92 31 39 31 52 32	.17 3 .93 3	0.24	29.53		23. 26. 30.	96 24 92 24 35 28	.49 .75	13.32	24.69 24.49 24.39 30.34 17.36
1877	ا۔۔ ا	56	57 51 51 60	56 0 00	57	57 5 57 5	8 57	59	60 57 62	61 4. 56 4.	9 4. 6 4.	8 5. 5 4.	5.6	5.1	4.9	4.9	5, I 4.5	5.1	5.I 4.4	5. I 4.3	5.0	18.	38 88	22.01 24.11	20.0 24.2 20.4 29.1	3 25. 1 25.	00 28 89 26	.61 2	8.95	26.49	34.4	26.	00 27	.50	5.68	28.87 25.65
1878 1879 1880 1881	63	56 64 57 62 54	57 51 58 58 52 59 55 57	559 58 58 58	57	57 5 57 5 57 5 57 5 60 6	58 58 54 60 60 60 57 57 57 57	54 61	53	63 4 53 4 63 5 60 5 58 5 56 4	3 4.	2 4. 8 4.	4.6	4.7	4.4	4.2	4.5	4.0	4.7	3.0	3,-	29.	13 1	24.20	20.4 29.1 22.8 36.2 28.9 28.9	7 20.	12 22	.30 2	9.35	26.93 31.25	25.13 32.18 26.58	27.	67 32	.86	32.68	27.77 32.97 38.65

A careful study of these curves will show many very interesting peculiarities. 1st, There is quite a marked minimum in the sun spot and magnetic declination curves between 1878 and 1879. 2d, The fluctuations in the latter correspond quite closely with those in the former, occurring, however, in many instances, from two to three months later. The enormous increase of sun spots in April, May, June, and July, 1885, has no such marked increase in the magnetic declination, and may be due to a large number of smaller spots having been counted in those months. 3d, There seems to be no close connection between the first two curves and either the range in air pressure or the mean temperature. 4th, There is a slight indication of a diminution of clouds and precipitation during the minimum of the sun spot period, but if there be a direct and important connection it is largely masked by other forces.

In presenting this method of investigation and these almost negative results,

In presenting this method of investigation and these almost negative results, it is not intended to assume that the question of a connection or non-connection is by any means settled. It is simply an indication that, in common with nearly all other investigations in this line, this method of attacking the problem gives only negative results. The meteorological elements are affected by so many causes, which serve to mask the real effects from any cosmical or supraterrestrial force, that we cannot hope to obtain a satisfactory solution until we are enabled to eliminate all other forces disturbing the one under consideration.

are enabled to eliminate all other forces disturbing the one under consideration.

A single example may be given in a possible line of investigation. It is evident that taking a monthly mean of any element will frequently smooth out the very effect we wish to study. Now, taking the temperature, it would seem that a daily curve of this element, when compared with the daily sun spot curve, might show a connection which could not otherwise be obtained, but before such comparison could be made it would be essential to examine the influence of day and night, high and low pressure, invisible vapor in the atmosphere, clouds, precipitation, etc. It is evident that clouds would have contrary effects at night and in the daytime; in the first case they would tend to increase the temperature, and in the latter to diminish it.

The whole problem is much more complicated than appears at first sight. Incidentally, several very interesting points are brought out by the last three curves. 1st, With very few exceptions the winters of odd years are relatively cold, and those of even years are warm. 2d, In general the warm months have the least clouds and the cool months the most. 3d, As was to be expected, the curves of cloudiness and precipitation are very similar

RESULTS OF ANEMOMETER OBSERVATIONS AT SEA. [By Prof. FRANK WALDO.]

On a voyage from Baltimore, United States, to Bremerhaven, Germany, from September 28th to October 15th, 1882, on board the steamship "Ohio," three anemometers were exposed—No. 527 on cross piece of the mainmast, about eighty feet above the water; No. 519 just over the front edge, and on the top of the awning frame of the bridge, about thirty-five feet above the water; No. 521 on the wheel-house, near the stern, about twenty-seven feet above the water. The anemometers were of the Robinson form, with cups 4 above the water. The anemometers were of the Robinson form, with cups 4 inches (101.6 mm.) in diameter and arms 6.72 inches (170.7 mm.). These are the dimensions of the standard instrument of the Signal Service. The anemometer on the bridge blew off early in the series of observations and was rendered unserviceable. The few readings are of use in showing results of the exposures. The instruments had all been compared with the Signal Service standard, from which they did not differ more than four per cent. -They were apparently new and in perfect condition at the time of mounting on ship-board, and, with the exception noted, remained in good condition as long as they were in use.

Analysis of the problem. We observe: 1st, the true direction of the ship's course; 2d, the motion of the ship in an hour; 3d, the estimated true direction of approach of the wind; 4th, the anemometer reading for the hour; this is the resultant effect of the velocity of the wind and of the steamer. Required the true velocity of the wind for the hour.

To avoid onerous calculation I have devised a graphical method of solving the triangle between the ship and the wind, as follows:

Lay off on a line on a convenient scale 0, 1, 2, 3, &c., parts which correspond to the anemometer reading. Lay off another line, beginning at 0 and forming an angle of 1 point (11½°) with this first line. On the second line mark off spaces 0, 1, 2, 3......16, 17, 18, corresponding to knots (sea miles); these to be on the same actual scale as the divisions of the first lines. Then through the points 1, 2, 3......16, 17, 18 draw lines parallel to the first line drawn. There points 1, 2, 8......16, 17, 18 draw lines parallel to the first line drawn. will be 18 of these lines.

The same is done for angles 2 points, 8 points, &c. Take a pair of dividers and use, as follows: We have given the anemometer reading, the space passed over by the steamer, and the angle, expressed in points, between the true reading. No record of hoisting the mainsail was kept, and this sail, though wind direction and the ship's course. Select the sheet on which the lines are small, might have deflected the wind, causing smaller readings on the mainsail was the course to probability of the ship o at the correct angle. Adjust the dividers to the number of miles recorded for | mast.

the hour by the anemometer, by means of the first line above. one leg of the dividers at the intersection of the two original lines. Sweep with the other leg until the free end falls on the parallel line drawn through the divisions corresponding to the space passed over by the ship in the hour. Let the free leg rest on this intersection and, by readjusting the dividers, let the other leg touch the end of the same parallel line, where it intersects the second line, drawn at the division which marks the space passed over by the ship. Take up the dividers unchanged and find how many miles on the first line drawn correspond to the distance between the points of the dividers. This distance is the true wind movement for the hour.

Table I shows: 1st, the wind estimated on the Beaufort scale, 0-12; 2d, the

number of hours for each anemometer corresponding to the wind velocity in the first column; 3d, the reduced mean anemometer indication for the same

time (true wind velocity):

TABLE I.

,		Hours	١.	Anemometer.							
Beaufort scale.	м. м.	w. н,	Bridge.	м. м.	w. п.	Bridge.					
1.5		3 3 21	4 4 1	2.5 11.0 12.3	6.3 9.3 13.1	14 I 18.8 18.7					
\$.5	19 1 21 37 37 19 12 5	12 24 1 18 36 34 11 8 3	1 18 18 4 2 1 1 2 2	13.4 14.1 23.1 19.9 27.5 27.2 24.1 28.4 32.4 46.2 47.2 51.8	10.0 16.4 22.3 19.4 30.5 32.6 37.7 39.3 40.5 43.7 44.8	23.9 19.5 19.1 29.2 26.8 35.9 30.6 43.4 43.4 346.8					

From Table I, by a graphical construction, we obtain, approximately, the wind velocity corresponding to each figure of the Beaufort scale at each exposure, as in Table II:

					7 V	BLL.			
1	2	3	4	5	6	7	8	9	Beaufort scale.
6	10	15	17	9Λ	97	30	49	54	Mainmast (80 feet).
7	10	13	17	23	32	40	48	47	Wheel-house (21 leet).
12	18	22	25	28	30	32	87	48	Bridge (35 feet).

Table III gives a comparison between these results and those of Prof. R. H. Scott, in England (Quarterly Journal Meteorological Society, Vol. II, p. 109). Column five is formed from Table II, by taking the mean of the three anemometers, giving a weight of 1 to the bridge and weights of 8 to the other two to allow for the number of observations:

' TABLE III.

Beaufort scale.	H. Scott.	F. W	aldo.	Mean of three anemometers.
	В. Н.	W. H.	м. м.	Мевп
7	8	,	6	7
2		10	13	ia
3	13 18		15	12 15 18 22
4	23	13	17	18
5	23 28	23	20	22
6	34	23 32	27	30
7	40	40	30	35
8	48	43	42	35 42
9	40 48 56	47	30 42 54	50

Comparing the results at twenty-seven and eighty feet, we find that up to 17 miles per hour there seems little difference, but from 17 to 42 the lower

